

## THE ARCIDAE OF THAILAND

By Vararin Vongpanich

Phuket Marine Biological Center, P.O. Box 60, Phuket 83000, Thailand.

### ABSTRACT

Arcid bivalves of Thailand belong to four genera: *Arca*, *Trisidos*, *Anadara* with two subgenera: *Scapharca* and *Tegillarca*, and *Barbatia*. Only the first three genera are included in this study. Eighteen species are recorded: *Arca navicularis* Bruguiere, *Arca ventricosa* Lamarck, *Arca avellana* Lamarck, *Trisidos tortuosa* (L.), *Trisidos semitorta* (Lamarck), *Anadara antiquata* (L.), *Anadara pilula* (Reeve), *Anadara troscheli* (Dunker), *Anadara ferruginea* (Reeve), *Anadara tricenicosta* (Nytz), *Anadara cuneata* (Reeve), *Anadara (Scapharca) mosambicana* (Bianconi), *Anadara indica* (Gmelin), *Anadara (Scapharca) inaequalis* (Bruguiere), *Anadara (Scapharca) vellicata* (Reeve), *Anadara clathrata* (Reeve), *Anadara (Tegillarca) granosa* (L.), *Anadara (Tegillarca) nodifera* (von Marten). The two species *Anadara mosambicana* and *Anadara cuneata* are new records for Thailand.