

Growth rate of *Trochus niloticus* L. fed on four species of benthic marine macroalgae

Eddy Soekendarsi, M.Iqbal Djawad & Yulianus Paongan



E. Soekendarsi, M.I. Djawad & Y. Paongan. 2001. Growth rate of *Trochus niloticus* L. fed on four species of benthic marine macroalgae. - Phuket Marine Biological Center Special Publication 25(1): 135-137.

T. niloticus were fed dry and fresh algae: *Caulerpa sertularoides* (Vahl) C. Agardh, *Ulva reticulata* Forsskal, *Padina australis* Hauck, and *Gracilaria salicornia* (C. Agardh) Dawson. Green macroalgae resulted in the best growth of *Trochus*, Brown macroalgae also contributed to an increase of the shell, but were inferior to green macroalgae.

Eddy Soekendarsi. Department of Biology, FMIPA - University of Hasanuddin, Makassar 90245, Indonesia.

M.Iqbal Djawad. Department of Fisheries, Faculty of Marine Science & Fisheries, University of Hasanuddin, Makassar 90245, Indonesia.

Yulianus Paongan. Department of Biology, FMIPA - University of Nusa Cendana, Kupang, Indonesia.

E-mail Eddy Soekendarsi: zad@indo.net.id