

A LIST OF POLYCHAETES (ANNELIDA) IN THE REFERENCE COLLECTION DATABASE OF THE PHUKET MARINE BIOLOGICAL CENTER, THAILAND

Charatsee Aungtonya, Santisuk Thaipal and Somchai Bussarawit

Phuket Marine Biological Center, P.O. Box 60, Phuket 83000, Thailand

ABSTRACT

A total of 1,101 polychaete records were registered at the Reference Collection of the Phuket Marine Biological Center in its computerized database as of 1 October 2001. These specimens were exclusively collected in Thai waters, the majority of them from the Andaman Sea. The collection contains 145 species representing 37 families, and of these 30 are holotypes and 28 are paratype lots. A list of these records, including PMBC catalogue numbers and locality information, is provided. The deposited materials will be valuable in further studies of polychaetes within the South-East Asian region, particularly the Andaman Sea.

INTRODUCTION

The Phuket Marine Biological Center (PMBC) was established in 1968 under the auspices of the Thai Department of Fisheries, Ministry of Agriculture, with financial support from the Danish Government. PMBC holds the largest collection of marine animals in Thailand, and is a significant site for taxonomic knowledge and biodiversity research.

The PMBC Reference Collection database was established and designed in 1997. Prior to this, hand-written registration cards were used. The database is in many ways superior in comparison with registration cards, *e.g.*, it is easy to search, information can easily be shared or updated and specimen catalogues can be printed (Bussarawit *et al.* 1999). The PMBC database presently holds over 14,000 records of marine species found in the south of Thailand.

The first checklist of polychaetes in the Reference Collection of the Phuket Marine Biological Center was published on the occasion of the 10th anniversary of the Center's opening. At that time there were only 4 species representing 3 families recorded from Thai waters (Nateewathana *et al.* 1981).

The polychaete specimens in the Reference Collection have mainly been collected during three projects: the Fifth Thai–Danish Expedition (1966), the Project on the Effect of Offshore Tin Mining on Benthic Communities (1980–83) and the Thai–Danish BIOSHELF surveys in the Andaman Sea off southwestern Thailand (1996–2000).

A list of the polychaetes from the Fifth Thai–Danish Expedition in the Andaman Sea was published by Phasuk (1992) and included 154 species in 39 families. However, not all specimens were deposited in the PMBC Reference Collection. The total of 807 accession numbers of polychaetes from this expedition deposited in the Collection contains only 115 species belonging to 31 families. The rest, which appears to contain several undescribed species (B. Phasuk, pers.comm.), was originally kept in Phasuk's private collection. Unfortunately, his ability to complete the study of this material was obstructed following his official retirement. It has recently (October 2001) been returned to the Reference Collection and will be catalogued following identification (see Hylleberg 2002, fig.1).

The material from the project on the effects of offshore tin mining on marine benthos on the west coast of Phuket has yielded a number of

polychaete species, including many new ones, which have been added to the PMBC Reference Collection (see Hylleberg and Nateewathana 1988, 1991a, 1991b, 1991c; Hylleberg *et al.* 1986; Nateewathana 1988, 1992; Nateewathana and Hylleberg 1986, 1991a, 1991b). However, the bulk of the polychaetes from this project is still unidentified and is kept at the Reference Collection, awaiting future study.

Much of the BIOSHELF polychaete material from 1996-1997 was studied during and following the Thai-Danida International Workshop on Polychaetes in 1997 (Aungtonya *et al.* 2000; this volume). The catalogue numbers of specimens in this material that have been deposited at the PMBC are not included in the table presented here but are to be found in the individual papers in this volume. The BIOSHELF polychaete material collected from 1998 to 2000 is currently being sorted and worked up. Selected groups will be distributed to various specialists.

The purpose of this paper is to provide a list of identified specimens of polychaetes in the Reference Collection. This material is useful as a reference source for the study of polychaetes in the South-East Asian region. Material can either be studied at the PMBC or requested on loan under the Center's regulations.

MATERIALS AND METHODS

The systematic arrangement of polychaetes in the PMBC Reference Collection was recently revised. Order names (previously Errantia and Sedentaria) and family names were changed in accordance with Fauchald (1977) and Fauchald and Rouse (1997). The table presented here is organized by family, listed alphabetically.

Information on each polychaete species, including PMBC catalogue numbers and the collection localities, was retrieved from the database. The existence of registered specimens in the collection was checked and missing specimens are noted in Table 1. Genera and species are listed alphabetically and retained as they appear in the original records. In Table 1, the locality of collection indicates only whether specimens were collected in the Andaman Sea (the southwest coast

of Thailand) or the Gulf of Thailand (the east coast and southeast coast of Thailand). Further details are available on request.

Dr. Danny Eibye-Jacobsen (Zoological Museum, University of Copenhagen, ZMUC) kindly confirmed the identity of specimens belonging to the families Glyceridae and Goniadidae. We have indicated his comments and identifications in the remarks column of Table 1 (see also Bggemann and Eibye-Jacobsen 2002).

RESULTS AND DISCUSSION

A total of 1,101 polychaete lots were deposited in the Reference Collection of the Phuket Marine Biological Center as of 1 October 2001 (Table 1). All specimens were collected in Thai waters, the majority of them in the Andaman Sea. A few additional records are from the Gulf of Thailand and one from Chao Phraya River. The collection contains 145 species, most of them named, representing 37 families. There are 58 type lots (30 holotypes and 28 paratype lots) in the collection.

Nateewathana *et al.* (1981) reported the presence of *Lepidonotus kumari* Rullier (Polynoidae), *Dendronereis arborifera* Peters and *Perinereis aibuhitensis* Grube (Nereididae), and *Parheteromastus tenuis* Monro (Capitellidae) (as *Paraheteromastus*) in the collection. However, for unknown reasons none of these species is presently represented by any registered material.

It is hoped that the present checklist will help draw the attention of specialists and students who are interested in studying polychaetes at the Reference Collection. In particular, a large number of unidentified polychaete specimens from previous collections needs to be worked up.

ACKNOWLEDGEMENTS

We express our appreciation to DANIDA for financial support of the Reference Collection Database Project, including the funding of computer sets and temporary staff during 1997-1999, under the Scientific Cooperation Programme between Denmark and Thailand (SCP). Sincere thanks are given to Mr. Praween Limpsaichol, the

Director of PMBC, and Dr. Jens Peter Thomson (the Chief Technical Advisor of SCP) for their encouragement in carrying out the Database Project. We thank Mrs. Ratchanee Sirivejabandhu, Mrs. Dongjan Srisuwan, and Dr. Andrew Davison for all their assistance during the project. Sincere

acknowledgement is given to Dr. Danny Eibye-Jacobsen (ZMUC) for kindly confirming the identity of specimens belonging to the families Glyceridae and Goniadidae and improving this manuscript, and to Mr. Ukkrit Satapoomin for his suggestions to a draft of this manuscript.

REFERENCES

- Aungtonya, C., S. Thaipal and O. Tendal. 2000. A preliminary report on the Thai–Danish BIOSHELF surveys (1996–2000) of the west coast of Thailand, Andaman Sea. *Phuket Marine Biological Center Research Bulletin* **63**: 53–76.
- Bggemann, M. and D. Eibye-Jacobsen. 2002. The Glyceridae and Goniadidae (Annelida: Polychaeta) of the BIOSHELF Project, Andaman Sea, Thailand. *Phuket Marine Biological Center Special Publication* **24**: 149–196.
- Bussarawit, S., A. Davison and C. Aungtonya. 1999. The design of the PMBC Reference Collection Database. *Phuket Marine Biological Center Special Publication* **19** (2): 407–411.
- Fauchald, K. 1977. The polychaete worms. Definitions and keys to the orders, families and genera. Allan Hancock Foundation, University of Southern California, Los Angeles. 188 pp.
- Fauchald, K. and G.W. Rouse. 1997. Polychaete systematics: Past and present. *Zoologica Scripta* **26**: 71–138.
- Hylleberg, J. 2002. The history of studies on the marine worms (Annelida: Polychaeta) of Thailand. *Phuket Marine Biological Center Special Publication* **24**: 13–16.
- Hylleberg, J. and A. Nateewathana. 1988. Polychaetes of Thailand. Nereididae (Part 2): *Ceratocephale* and *Gymnonereis*, with description of two new species and notes on the subfamily Gymnonereidinae. *Phuket Marine Biological Center Research Bulletin* **49**: 1–20.
- Hylleberg, J. and A. Nateewathana. 1991a. Temporal and spatial distribution of subtidal magelonid polychaetes at Phuket Island, Andaman Sea, Thailand. *Ophelia Supplement* **5**: 573–578.
- Hylleberg, J. and A. Nateewathana. 1991b. Spatial and temporal distribution of spionid polychaetes at Phuket Island, the Andaman Sea. *Bulletin of Marine Science* **48**: 346–357.
- Hylleberg, J. and A. Nateewathana. 1991c. Polychaetes of Thailand. Spionidae (Part 1): *Prionospio* of the *steenstrupi* group with description of eight new species from the Andaman Sea. *Phuket Marine Biological Center Research Bulletin* **55**: 1–32.
- Hylleberg, J., A. Nateewathana and S. Bussarawit. 1986. Polychaetes of Thailand. Nereididae (Part 1): *Perinereis* and *Pseudonereis* with notes on species of commercial value. *Phuket Marine Biological Center Research Bulletin* **43**: 1–22.
- Nateewathana, A. 1988. *Heptaceras hyllebergi* n. sp. (Polychaeta: Onuphidae) from Kata Beach, west coast of Phuket Island, Andaman Sea, Thailand, with notes on the habitat. *Phuket Marine Biological Center Research Bulletin* **48**: 1–9.
- Nateewathana, A. 1992. Polychaetes of Thailand, Nereididae (Part 3): *Solomononereis phuketensis* n. sp. from euhaline environments in the Andaman Sea, Thailand. *Phuket Marine Biological Center Research Bulletin* **57**: 89–96.
- Nateewathana, A. and J. Hylleberg. 1986. Nephtyid polychaetes from the west coast of Phuket Island, Andaman Sea, Thailand, with description of five new species. *Proceedings of the Linnean Society of New South Wales* **108**: 195–215.
- Nateewathana, A. and J. Hylleberg. 1991a. Magelonid polychaetes in the Andaman Sea, Thailand, with description of eight new species. *Ophelia Supplement* **5**: 169–184.

- Nateewathana, A. and J. Hylleberg. 1991b. Characters used to separate species of *Prionospio* (Spionidae) of the *steenstrupi* group from the Andaman Sea, Thailand. *Bulletin of Marine Science* **48**: 277–279.
- Nateewathana, A., P. Tantichodok, S. Bussarawit and R. Sirivejabandhu. 1981. Marine organisms in the Reference Collection. *Phuket Marine Biological Center Research Bulletin* **28**: 43–86.
- Phasuk, B. 1992. Preliminary report on the polychaetes from the Fifth Thai–Danish Expedition along the Andaman Sea coast of Thailand. *Phuket Marine Biological Center Research Bulletin* **57**: 77–87.

Table 1 List of polychaete lots in the Reference Collection database of the Phuket Marine Biological Center, Thailand. Families are listed in alphabetical order. AS = Andaman Sea, CP = Chao Phraya River, GT = Gulf of Thailand.

Family	Species	PMBC cat. no.	Locality	Remarks
ACOETIDAE	<i>Eupanthalis oculata</i> Hartman, 1944	12791–2	AS	
	<i>Panthalis marginata</i> Hartman, 1939	12905–15	AS	
	<i>Polyodontes</i> sp.	12973–4, 13629	AS	
AMPHARETIDAE	<i>Amphicteis</i> cf. <i>gunneri malayensis</i> Caullery, 1944	12980	AS	
	<i>Auchenoplax crinita</i> Ehlers, 1887	12133–40	AS	
	<i>Eclysippe vanelli</i> (Fauvel, 1936)	12675–7	AS	
	<i>Eclysippe</i> sp.	12689–93	AS	
	<i>Samytha</i> sp.	12711–3	AS	
AMPHINOMIDAE	<i>Chloeia amphora</i> Horst, 1910	13007–8	AS	
	<i>Chloeia flava pulchella</i> (Baird, 1870)	6023	GT	
	<i>Chloeia parva</i> Baird, 1870	13009	AS	
	<i>Chloeia violacea</i> Horst, 1910	12962–72, 12981–96, 13700–6	AS	
	<i>Hipponoa</i> sp.	13595–97	AS	
CAPITELLIDAE	<i>Capitellethus branchiferus</i> Gallardo, 1968	12790	AS	
	<i>Capitomastus</i> cf. <i>minimus</i> (Langerhans, 1881)	12997–9	AS	
	<i>Dasybranchus lumbricoides</i> Grube, 1878	13005–6	AS	
	<i>Mastobranthus</i> sp. A	12124–9	AS	
	<i>Notomastus latericeus</i> Sars, 1851	12849–74, 13630–2	AS	
	<i>Scyphoproctus gravieri</i> Okuda, 1940	13010	AS	
	<i>Scyphoproctus steinitzii</i> Day, 1965	13011–13	AS	
CHAETOPTERIDAE	<i>Chaetopterus variopedatus</i> Renier, 1803	6019	AS	
	<i>Mesochaetopterus japonicus</i> Fujirara, 1934	12346–57, 12375	AS	

Table 1 (continued).

Family	Species	PMBC cat. no.	Locality	Remarks
	<i>Phyllochaetopterus claparedii</i> McIntosh, 1885	12378–81	AS	
CIRRATULIDAE	<i>Cirriformia</i> sp.	13606–8	AS	
DORVILLEIDAE	<i>Microdorvillea phuketensis</i> Westheide, 1992	7270	AS	HOLOTYPE
EULEPETHIDAE	<i>Pareulepis malayana</i> (Horst, 1913)	12951	AS	
EUNICIDAE	<i>Eunice indica</i> Kinberg, 1865	12568–75, 12582–5, 13613–5	AS	
	<i>Marphysa adenensis</i> Gravier, 1900	12626–34	AS	
FLABELLIGERIDAE	<i>Pherusa bengalensis</i> (Fauvel, 1932)	12181–84	AS	
	<i>Pherusa curvisetis</i> (Caullery, 1944)	11988–90	AS	
	<i>Pherusa eruca indica</i> (Fauvel, 1928)	11986–7	AS	
GLYCERIDAE	<i>Glycera alba</i> (Müller, 1788)	12646 12647–8, 13616	AS AS	confirmed
	<i>Glycera chirori</i> Izuka, 1912	13609	AS	<i>G. nicobarica</i> Grube, 1868
		13610	AS	? <i>G. brevicirris</i> Grube, 1870
	<i>Glycera gigantea</i> Quatrefages, 1865	12597, 12599 12598	AS AS	? <i>G. nicobarica</i> Grube, 1868 <i>G. onomichiensis</i> Izuka, 1912
		13612	AS	Polynoidae sp.
	<i>Glycera prashadi</i> Fauvel, 1932	13618	AS	<i>G. onomichiensis</i> Izuka, 1912
	<i>Glycera tessellata</i> Grube, 1863	12541, 12543–4, 13617 12542	AS AS	<i>G. oxycephala</i> Ehlers, 1887 <i>G. nicobarica</i> Grube, 1868
		12545	AS	<i>G. alba</i> (Müller, 1776)

Table 1 (continued).

Family	Species	PMBC cat. no.	Locality	Remarks
GONIADIDAE	<i>Goniada emerita</i> Audouin and Milne Edwards, 1833	12453	AS	<i>Goniada</i> sp. (not <i>G. emerita</i>)
	<i>Goniada littorea</i> Hartman, 1950	13699	AS	<i>Goniada</i> cf. <i>littorea</i> Hartman, 1950
	<i>Goniada multidentata</i> Arwidsson, 1899	12559–63 12564	AS AS	confirmed <i>Goniada</i> n. sp.
HESIONIDAE	<i>Hesione</i> sp.	12141	AS	
	<i>Hesionides bengalensis</i> Westheide, 1992	7269	AS	HOLOTYPE
	<i>Leocrates indicus</i> Horst, 1921	12565	AS	
LUMBRINERIDAE	<i>Lumbrineris latreilli</i> Audouin and Milne Edwards, 1834	12920–45, 13593–4	AS	
	<i>Lumbrineris albidentata</i> Ehlers, 1908	13709	AS	
MAGELONIDAE	<i>Magelona alleni</i> Wilson, 1958	12467–76, 12885, 12916–9	AS	
	<i>Magelona californica</i> Hartman, 1944	13697–8	AS	
	<i>Magelona cincta</i> Ehlers, 1908	4346–50	AS	
	<i>Magelona crenulifrons</i> Gallardo, 1968	4292–308, 13676–84	AS	
	<i>Magelona kamala</i> Nateewathana and Hylleberg, 1991	4283	AS	HOLOTYPE
		4284	AS	PARATYPE
		4285–7	AS	
	<i>Magelona lenticulata</i> Gallardo, 1968	13648–9	AS	
	<i>Magelona methae</i> Nateewathana and Hylleberg, 1991	4344	AS	HOLOTYPE
		4345	AS	PARATYPE
	<i>Magelona mickminni</i> Nateewathana and Hylleberg, 1991	3174	AS	HOLOTYPE
		3175	AS	PARATYPE
		4356–64	AS	
	<i>Magelona noppi</i> Nateewathana and Hylleberg, 1991	4280	AS	HOLOTYPE
		4281	AS	PARATYPE
		4282	AS	
	<i>Magelona papillicornis</i> Møller, 1858	13647	AS	
<i>Magelona pectinata</i> Nateewathana and Hylleberg, 1991	3165	AS	HOLOTYPE	
	3166	AS	PARATYPE	
	3167–8, 3176–8, 4169–75	AS		
<i>Magelona peterseni</i> Nateewathana and Hylleberg, 1991	4352	AS	HOLOTYPE	
	4351	AS	PARATYPE	
	4353–5	AS		

Table 1 (continued).

Family	Species	PMBC cat. no.	Locality	Remarks
	<i>Magelona pygmaea</i> Nateewathana and Hylleberg, 1991	4219 4220 4221–48	AS AS AS	HOLOTYPE PARATYPE
	<i>Magelona tinae</i> Nateewathana and Hylleberg, 1991	3179 3180 4249–71	AS AS AS	HOLOTYPE PARATYPE
MALDANIDAE	<i>Asychis gangeticus</i> Fauvel, 1932	12193–203	AS	
	<i>Axiothella obockensis</i> (Gravier, 1905)	3792, 12149–57	AS	
	<i>Boguea panwaensis</i> Meyer and Westheide, 1997	13577	AS	HOLOTYPE
	<i>Euclymene annandalei</i> Southern, 1921	12143–48, 12225, 12230, 12701–5	AS	
	<i>Euclymene</i> cf. <i>santanderensis</i> (Rioja, 1917)	12698–700	AS	
	<i>Maldane sarsi</i> Malmgren, 1865	12226	AS	
NEPHTYIDAE	<i>Aglaophamus dibranchus</i> (Grube, 1877)	12581	AS	
	<i>Aglaophamus dicirrus</i> Hartman, 1950	12566, 12837–48	AS	
	<i>Aglaophamus phuketensis</i> Nateewathana and Hylleberg, 1986	3159 3160 5111	AS AS AS	HOLOTYPE PARATYPE
	<i>Aglaophamus sinensis</i> (Fauvel, 1932)	12576–8	AS	
	<i>Aglaophamus sphaerocirrata</i> (Wesenberg-Lund, 1949)	12744–6	AS	
	<i>Aglaophamus urupani</i> Nateewathana and Hylleberg, 1986	3102 3103 5110	AS AS AS	HOLOTYPE PARATYPE
	<i>Aglaophamus</i> cf. <i>verrilli</i> (McIntosh, 1885)	3169, 5116	AS	
	<i>Inermonephtys</i> cf. <i>gallardi</i> Fauchald, 1968	3161–2, 5112	AS	
	<i>Inermonephtys patongi</i> Nateewathana and Hylleberg, 1986	3170 3171 5114	AS AS AS	HOLOTYPE PARATYPE
	<i>Micronephtys sphaerocirrata</i> (Wesenberg-Lund, 1949)	5109	AS	
	<i>Nephtys danida</i> Nateewathana and Hylleberg, 1986	3163 3164 5113	AS AS AS	HOLOTYPE PARATYPE

Table 1 (continued).

Family	Species	PMBC cat. no.	Locality	Remarks
	<i>Nephtys phasuki</i> Nateewathana and Hylleberg, 1986	3172 3173	AS AS	HOLOTYPE PARATYPE
	<i>Nephtys</i> sp.	5115 12747-64	AS AS	
NEREIDIDAE	<i>Ceratocephale andaman</i> Hylleberg and Nateewathana, 1988	6016 6017	AS AS	HOLOTYPE PARATYPE
		6003-7	AS	
	<i>Ceratocephale fauveli</i> Pillai, 1961	12740	AS	
	<i>Gymnonereis</i> cf. <i>fauveli</i> (Pillai, 1961)	6008, 6011, 6013, 6015	AS	
		6009, 6012, 6014	GT	
	<i>Gymnonereis phuketensis</i> Hylleberg and Nateewathana, 1988	6000 6001	AS AS	HOLOTYPE PARATYPE
		6002, 6010	AS	
	<i>Namalycastis abiuma</i> (Müller in Grube, 1872)	4615 4613-4	AS GT	
	<i>Namalycastis fauveli</i> Nageswara Rao, 1981	7244-5	-	
	<i>Namalycastis indica</i> (Southern, 1921)	4616-7, 4619	GT	
		4618	CP	
	<i>Perinereis aibuhitensis</i> Grube, 1878	3865-8	AS	
	<i>Perinereis quatrefagesi</i> (Grube, 1878)	3946-9, 3954	AS	
		3950-2	GT	
	<i>Perinereis</i> cf. <i>rhombodonta</i> Wu, Sun and Yang, 1981	10091	GT	
	<i>Perinereis singaporiensis</i> (Grube, 1878)	3942, 3953	AS	
	<i>Perinereis striolata</i> Grube, 1878	3943-5	AS	
	<i>Pseudonereis anomala</i> Gravier, 1901	3848-50	AS	
	<i>Pseudonereis gallapagensis</i> Kinberg, 1866	3851	AS	
	<i>Solomononereis phuketensis</i> Nateewathana, 1992	8743 8744	AS AS	HOLOTYPE PARATYPE
		8745	AS	

Table 1 (continued).

Family	Species	PMBC cat. no.	Locality	Remarks	
ONUPHIDAE	<i>Heptaceras hyllebergi</i> Nateewathana, 1988	4696	AS	HOLOTYPE	
		4697	AS	PARATYPE	
		4698–700, 10666–9	AS		
		<i>Hyalinoecia</i> sp.	12678–84	AS	
		<i>Nothria holobranchiata</i> (Marenzeller, 1879)	12820–36	AS	
		<i>Onuphis eremita</i> Audouin and Milne Edwards, 1833	12232–43, 12245–53	AS	
		<i>Onuphis</i> sp.	12586–88	AS	
OPHELIIDAE	<i>Ammotrypane grandis</i> Pillai, 1961	12978	AS		
	<i>Armandia longicaudata</i> (Cauullery, 1944)	12376–7	AS		
	<i>Travisia</i> cf. <i>japonica</i> Fujiwara, 1933	12393,	AS		
		12397, 12399, 12403–4			
ORBINIIDAE	<i>Haploscoloplos elongatus</i> (Johnson, 1901)	12382–7	AS		
	<i>Orbinia vietnamensis</i> Gallardo, 1968	12339–45	AS		
	<i>Scoloplos (Leodamas) gracilis</i> Pillai, 1961	12600–25	AS		
	<i>Scoloplos (Leodamas) rubra orientalis</i> Gallardo, 1968	12975	AS		
	<i>Scoloplos marsupialis</i> Southern, 1921	12706–10	AS		
OWENIIDAE	<i>Owenia fusiformis</i> (Delle Chiaje, 1844)	12454–66	AS		
PECTINARIIDAE	<i>Pectinaria</i> sp.	12635–45	AS		
PILARGIDAE	<i>Ancistrosyllis albini</i> Langerhans, 1881	12765–8	AS		
	<i>Sigambra phuketensis</i> Licher and Westheide, 1997	9211	AS	HOLOTYPE	
		9212–3	AS	PARATYPES	
		13748	AS		
PISIONIDAE	<i>Pisionidens indica</i> (Aiyar and Alikunhi, 1940)	10682–702	AS		
POECILOCHAETIDAE	<i>Poecilochaetus paratropicus</i> Gallardo, 1968	12209–23	AS		
POLYNOIDAE	<i>Iphione muncata</i> (Savigny, 1818)	6022	AS		

Table 1 (continued).

Family	Species	PMBC cat. no.	Locality	Remarks
SABELLARIIDAE	<i>Lygdamis nesiotus</i> (Chamberlin, 1919)	12696–7	AS	
SABELLIDAE	<i>Branchiomma cingulata</i> Grube, 1870	12026	AS	
	<i>Branchiomma serratibranchis</i> Grube, 1878	12004–11, 12013–4, 12046–9, 12080–1, 12102	AS	
	<i>Chone infundibuliformis</i> Krøyer, 1856	12082–9	AS	
	<i>Chone trilobata</i> Gallardo, 1968	12043–5	AS	
	<i>Chone</i> sp.	12038–42	AS	
	<i>Megalomma pacifica</i> Johansson, 1927	12020–5	AS	
SERPULIDAE	<i>Ditrupa arietina</i> (Müller, 1776)	12373	AS	
SIGALIONIDAE	<i>Psammolyce antipoda</i> (Schmarda, 1861)	12977	AS	
	<i>Psammolyce malayana</i> Horst, 1913	13626, 13685–8	AS	
	<i>Sigalion capense</i> Day, 1960	13017, 13622	AS	
	<i>Sthenelais articulata</i> Kinberg, 1855	13018–20, 13624	AS	
	<i>Sthenelais orientalis</i> Potts, 1910	13646	AS	
	<i>Sthenelais zeylanica</i> Willey, 1905	13588–92, 13627–8	AS	
	<i>Sthenelanella uniformis</i> Moore, 1910	13625, 13643–5, 13670–5	AS	
	<i>Sthenolepis yhleni</i> (Malmgren, 1867)	13689–96	AS	
	<i>Thalenessa lewisii</i> (Berkeley and Berkeley, 1939)	12875–84, 13633–42	AS	
	<i>Thalenessa</i> sp.	13623	AS	
SPIONIDAE	<i>Polydora caeca</i> (Cuvier, 1843)	13603–5	AS	
	<i>Prionospio andamanensis</i> Hylleberg and Nateewathana, 1991	7211, 7212, 7213, 7241–3	AS AS AS	HOLOTYPE PARATYPE
	<i>Prionospio cornuta</i> Hylleberg and Nateewathana, 1991	7214, 7215, 7216–9, 7234, 7235, 7266	AS AS AS AS AS	HOLOTYPE PARATYPE PARATYPE

Table 1 (continued).

Family	Species	PMBC cat. no.	Locality	Remarks
	<i>Prionospio ehlersi</i> Fauvel, 1928	12366–9	AS	
	<i>Prionospio henriki</i> Hylleberg and Nateewathana, 1991	7220 7221	AS AS	HOLOTYPE PARATYPE
		7222	AS	
	<i>Prionospio komaeti</i> Hylleberg and Nateewathana, 1991	7226 7227	AS AS	HOLOTYPE PARATYPE
		7228	AS	
	<i>Prionospio malayensis</i> Caullery, 1914	12438–51, 12490–540, 12732–7	AS	
	<i>Prionospio cf. malayensis</i> Caullery, 1914	7223–5, 7229, 7265	AS	
	<i>Prionospio neenae</i> Hylleberg and Nateewathana, 1991	7230 7231	AS AS	HOLOTYPE PARATYPE
		7232	AS	
	<i>Prionospio nielseni</i> Hylleberg and Nateewathana, 1991	7233	AS	HOLOTYPE
	<i>Prionospio phuketensis</i> Hylleberg and Nateewathana, 1991	7236 7237	AS AS	HOLOTYPE PARATYPE
		7267	AS	
	<i>Prionospio pinnata</i> Ehlers, 1901	12276–300	AS	
	<i>Prionospio runei</i> Hylleberg and Nateewathana, 1991	7238 7239	AS AS	HOLOTYPE PARATYPE
		7240	AS	
	<i>Prionospio tenuis</i> (Verrill, 1880)	12694–5	AS	
	<i>Pseudopolydora antennata</i> (Clap- ar de, 1870)	12979	AS	
	<i>Pseudopolydora</i> sp.	12593–6	AS	
	<i>Scolecopsis (Scolecopsis) laciniata</i> Eibye-Jacobsen, 1997	11665	AS	HOLOTYPE, lost during shipment
		11666–72	AS	PARATYPES, lost during shipment
		11673–86	AS	lost during shipment
		17524	AS	PARATYPE, originally ZMUC-POL- 433 (to replace lost HOLO- TYPE PMBC 11665)

Table 1 (continued).

Family	Species	PMBC cat. no.	Locality	Remarks
	<i>Spio</i> sp.	12477–89	AS	
	<i>Spiophanes bombyx</i> (Clapar de, 1870)	12685–8	AS	
	<i>Spiophanes</i> sp.	13582–3	AS	
STERNASPIDAE	<i>Sternaspis scutata</i> (Renier, 1807)	11984, 13720	AS	
SYLLIDAE	<i>Sphaerosyllis hermaphrodita</i> West- heide, 1990	6929	AS	HOLOTYPE
TEREBELLIDAE	<i>Amaeana occidentalis</i> Hartman, 1944	11981–3	AS	
	<i>Amaeana trilobata</i> (Sars, 1863)	11994–2000	AS	
	<i>Amaeana</i> sp.	12204–8, 13578	AS	
	<i>Loimia medusa</i> (Savigny, 1818)	12027–37	AS	
	<i>Loimia medusa annulifilis</i> (Grube, 1878)	11985, 12110–1	AS	
	<i>Pista fasciata</i> (Grube, 1870)	11980	AS	
	<i>Pista typha</i> Grube, 1878	12256, 12258, 12267, 12301–13	AS	
	<i>Streblosoma</i> cf. <i>amboinense</i> Caul- lery, 1944	11991–3	AS	
	<i>Streblosoma</i> sp.	12331–8	AS	
	<i>Thelepus robustus</i> Grube, 1878	12090–5	AS	
TRICHOBRANCHIDAE	<i>Terebellides stroemi</i> Sars, 1835	12413–37, 12668–74	AS	
TROCHOCHAETIDAE	<i>Trochochaeta orissae</i> (Fauvel, 1932)	6018	AS	